

A REMARK ON VIRTUAL ORIENTATIONS FOR COMPLETE INTERSECTIONS

VADIM SCHECHTMAN

The aim of this note is to give a simple definition of genus zero *virtual orientation classes* (or *fundamental classes*) for projective complete intersections or, more generally, complete intersections in convex varieties, and to prove a *push forward formula* (see Lemmas 1 and 2 below) for them.

1. Let X be a smooth complex projective variety. Let X_β denote the stack of stable maps $f : C \rightarrow X$ of genus g curves with n marked points, such that $f_*([C]) = \beta \in H_2(X)$. Here n and g will be fixed and suppressed from the notations. Let $\pi_\beta : \tilde{X}_\beta \rightarrow X_\beta$ be the universal curve, and $\psi_\beta : \tilde{X}_\beta \rightarrow X$ be the canonical map.

A vector bundle N over X is called *convex* if $R^1\pi_{\beta*}\psi_\beta^*N = 0$ for all β . In this case we will use the notation N_β for the vector bundle $\pi_{\beta*}\psi_\beta^*N$ over X_β . The variety X is called convex if the tangent bundle $\mathcal{T}_X$ is convex.

Example. Let $X = \mathbb{P}^N$, and $g = 0$. Then X is convex, and any bundle of the form $N = \bigoplus \mathcal{O}(l_a)$ ($l_a \geq 0$) is convex.

2. Let X be a convex variety and N a convex bundle over X . Let $s : X \rightarrow N$ be a *regular* (i.e. transversal to the zero section) section of N , $i : Y := s^{-1}(0) \hookrightarrow X$ the subscheme of its zeros.

Set $Y_\beta := \coprod Y_\gamma$, the disjoint union over all $\gamma \in H_2(Y)$ mapping to β . The section s induces the section $s_\beta : Y_\beta \rightarrow N_\beta$, and we have $Y_\beta = s_\beta^{-1}(0)$. We denote by i_β the embedding $Y_\beta \hookrightarrow X_\beta$.

Definition 1. The *virtual dimension* $\dim^{virt}(Y_\beta)$ is the number $\dim(X_\beta) - \text{rk}(N_\beta)$.

The section s is not necessarily regular. If it is, then the virtual dimension of Y_β coincides with its usual dimension.

3. Construction. Our goal is to define certain class in the homology Chow group $A_{\dim^{virt}(Y_\beta)}(Y_\beta)$. We will use the key construction from the intersection theory, [F], 6.1. We have a cartesian square

$$\begin{array}{ccc} Y_\beta & \xrightarrow{i_\beta} & X_\beta \\ i_\beta \downarrow & & \downarrow s_d \\ X_\beta & \xrightarrow{s_\beta} & N_\beta \end{array}$$

where s_0 denotes the zero section. Let C be the normal cone of Y_β in X_β . We have the canonical closed embedding $C \hookrightarrow i_\beta^*N_\beta$, whence the class $[C] \in A_d(i_\beta^*N)$ where $d =$

$\dim(C) = \dim(X_\beta)$. We set by definition

$$[Y_\beta]^{virt} := (p^*)^{-1}([C]).$$

Here $p^* : A_*(Y_\beta) \xrightarrow{\sim} A_{*+\text{rk}(N)}(i_\beta^* N)$ is the isomorphism induced by the projection $p : i_\beta^* N \longrightarrow Y_\beta$.

4. Push forward formula. Lemma 1. *We have*

$$i_{\beta*}([Y_\beta]^{virt}) = [X_\beta] \cap e(N_\beta)$$

where e denotes the Euler (top Chern) class.

This formula was written down as a conjecture by Yu.I.Manin (talk at MPI, February 1997).

Proof. We can identify $s_0 : X_\beta \longrightarrow N_\beta$ with the embedding of the scheme of zeros of the regular *diagonal* section of the vector bundle $N_\beta \times N_\beta \longrightarrow N_\beta$, and then apply [F], 6.3.4. $\square$

5. Iteration. Let M be a convex vector bundle over Y , and $j : Z \hookrightarrow Y$ be the subscheme of zeros of a regular section $t : Y \longrightarrow M$. We set by definition,

$$\dim^{virt}(Z_\beta) := \dim^{virt}(Y_\beta) - \text{rk}(M_\beta).$$

We define

$$[Z_\beta]^{virt} \in A_{\dim^{virt}(Z_\beta)}(Z_\beta)$$

by the same construction as in no. 3, with $[X_\beta]$ replaced by $[Y_\beta]^{virt}$.

If $M = i^* N'$ and t is the restriction of a regular section $s' : X \longrightarrow N'$ then Z may be defined as the subscheme of zeros of the regular section $s \oplus s' : X \longrightarrow N \oplus N'$, and the definition of $[Z_\beta]^{virt}$ at one step using $s \oplus s'$ coincides with the previous one (which used first s , then t).

We have a generalization of Lemma 1:

6. Push forward formula. II. Lemma 2. *We have*

$$j_{\beta*}([Z_\beta]^{virt}) = [Y_\beta]^{virt} \cap e(M_\beta).$$

The proof is the same as in Lemma 1.

7. The above constructions have obvious equivariant versions.

I am grateful to A.Goncharov and especially to Yu.I.Manin, for very useful discussions.

REFERENCES

- [F] W.Fulton, Intersection theory, Springer-Verlag, Berlin et al., 1984.

MAX-PLANCK-INSTITUT FÜR MATHEMATIK, GOTTFRIED-CLAREN-STRASSE 26, 53225 BONN, GERMANY

E-mail address: `vadik@mpim-bonn.mpg.edu`